

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
27 May 2004 (27.05.2004)

PCT

(10) International Publication Number
WO 2004/043898 A1

(51) International Patent Classification⁷: **C07C 201/02,**
203/04, C01B 21/46

CASTALDI, Graziano [IT/IT]; Via Livia Gallina, 5,
I-28072 Briona (IT).

(21) International Application Number:
PCT/EP2003/012376

(74) Agents: **MINOJA, Fabrizio** et al.; Bianchetti Bracco Mi-
noja S.r.l., Via Rossini, 8, I-20122 Milano (IT).

(22) International Filing Date:
6 November 2003 (06.11.2003)

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR,
CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD,
GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR,
KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN,
MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU,
SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA,
UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
MI2002A 002410
14 November 2002 (14.11.2002) IT

(84) Designated States (*regional*): ARIPO patent (BW, GH,
GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE,
SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA,
GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): **DI-
NAMITE DIPHARMA S.P.A.** (In Abbreviated form
DIPHARMA S.P.A.) [IT/IT]; Via XXV Maggio, 40,
I-33036 Mereto di Tomba (IT).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **FRANCESCUTTI,**
Nevio [IT/IT]; Via Caravaggio, 21, I-33070 San Giovanni
di Casarsa (IT). **SCUBLA, Tiziano** [IT/IT]; Via Bo-
nanni, 32, I-33037 Pasian di Prato (IT). **GORASSINI,**
Fausto [IT/IT]; Via Codroipo, 30, I-33100 Udine (IT).

Published:

— with international search report

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

(54) Title: A PROCESS FOR THE MONONITRATION OF ALKANEDIOLS


(57) Abstract: A process for the preparation of compounds of formula: HO-A-ONO₂ (I) wherein A is a C₂-C₆ alkylene chain by nitration of the corresponding alkanediols with "stabilised" nitric acid is herein disclosed. The process is safer to operators and allows to obtain advantageous yields on an industrial scale.

A PROCESS FOR THE MONONITRATION OF ALKANEDIOLS

FIELD OF THE INVENTION

The present invention provides an efficient process for the mononitration of alkanediols, which is easily controllable and safe to operators.

In particular, the invention provides a process for industrial scale production of compounds of formula



wherein A is a straight or branched C₂-C₆ alkylene chain.

BACKGROUND OF THE INVENTION

The compounds of formula (I) are useful intermediates for the synthesis of NO-releasing non steroidal anti inflammatory drugs ("NO-NSAIDs"). Examples of NO-NSAIDs are NO-acetylsalicylic acid, NO-diclofenac, NO-naproxen, NO-ketoprofen and NO-ibuprofen. Like conventional non steroidal anti inflammatory drugs, NO-NSAIDs have antipyretic and anti-inflammatory activity, but show reduced gastrointestinal toxicity and are endowed with further useful pharmacological properties. WO 98/25918 and WO 01/10814 in particular disclose the use of compounds of formula (I) in the preparation of NO-NSAIDs.

The production of mononitrated alkanediols on an industrial scale through reduction or hydrolysis of dinitrated alkanediols is poorly selective and troublesome, as the synthesis of dinitrated alkanediols involves problems during production, storage and transport similar to those encountered with nitroglycerine.

It is well known that the mononitration of alcohols (for example glycerin) can be usually carried out with concentrated nitric acid, optionally in

the presence of sulfuric acid and urea to remove nitrous acid from the reaction mixture. However, this reaction is poorly selective and quickly leads to formation of dinitrated alkanediols due to the fact that the kinetics is not easy to control. Moreover by carrying out the same reaction on short chain alkanediols strong decomposition reactions are easily triggered. WO 98/25918 also discloses alkanediols mononitration with silver nitrate or fuming nitric acid as nitrating agents, in particular with concentrated nitric acid and acetic anhydride. The mononitration of alkanediols of formula (I) with the above methods is very poorly selective and troublesome on an industrial scale. In particular, dangerous decompositions easily occur, since a strong oxidizing agent (nitric acid) is contacted with extremely unstable substrates, in particular with alkanediols wherein A is a C₂-C₄ alkylene chain.

There is therefore the need to develop a more selective and safer industrial process for the mononitration of diol compounds which also allows to obtain compounds of formula (I) in good yields.


DETAILED DESCRIPTION OF THE INVENTION

It has now been found that it is possible to selectively carry out the mononitration of alkanediols in high yield and under easily controllable conditions using "stabilised" nitric acid.

Object of the present invention is therefore a process for the preparation of a compound of formula:



wherein A is a C₂-C₆ alkylene chain,
comprising the nitration of a compound of formula



wherein A is as defined above,
with "stabilised" nitric acid.

Preferred compounds of formula (I) obtainable with the process of the invention are ethanediol-mononitrate; 1,3-propanediol-mononitrate; 1,4-butanediol-mononitrate; 1,5-pentanediol-mononitrate or 1,6-hexanediol-mononitrate, more preferably 1,4-butanediol-mononitrate.

As used herein, the term "stabilised nitric acid" means a specific preparation of nitric acid, which is a further object of the invention. This preparation consists of nitric acid diluted in water to a concentration of about 83 - 85%, preferably about 84.5 - 84.8% w/w, and is substantially free from nitrous acid and nitrogen oxides. The expression "substantially free from nitrous acid and nitrogen oxides" means that the concentration of nitrous acid and nitrogen oxides is typically lower than 10 p.p.m., preferably lower than 5 p.p.m.

Stabilization of nitric acid can be achieved by a process which comprises dilution of fuming nitric acid with water and treatment of the diluted nitric acid with an agent able to remove nitrous acid and nitrogen oxides (hereinafter referred to as "(the) agent/s"). It will be appreciated that the same result can also be obtained by adding an aqueous solution of the agent to fuming nitric acid.

Examples of agents are urea and sulfamic acid, preferably urea.

The contact time of the agents with nitric acid and their amount should be sufficient to make the nitric acid substantially free from the nitrous acid and nitrogen oxides, but should not salify an excessive amount of nitric acid, as this would lower the titre and the nitrating power. The absence of nitrogen oxides and nitrous acid in the "stabilised" nitric acid can be evaluated, for example, by visual inspection (colour), as well as by permanganate titration.

"Stabilised" nitric acid is prepared by adding 83 - 85% w/w nitric acid (which usually contains from about 0.06 to about 0.12% of nitrous acid and nitrogen oxides) with an amount of agent ranging from about 0.3 to about 1%

w/w. To completely remove nitrous acid and nitrogen oxides the time contact with nitric acid ranges from about 80 minutes to about 130 minutes.

In the case of urea, the amount typically ranges from about 0.6 to about 1% w/w, preferably from about 0.7 to about 1% w/w and the time contact is preferably from about 95 to about 120 minutes.

The "stabilised" nitric acid according to the invention must be used within about three hours from stabilization, as relevant amounts of nitrous acid and nitrogen oxides are released again over time.

The mononitration of compounds of formula (II) with "stabilised" nitric acid is preferably carried out in a water-immiscible chlorinated organic solvent, hereinafter referred to as "chlorinated solvent". Examples of chlorinated solvents are mono-, di-, tri- and tetra-chloro C_1 - C_4 alkyl hydrocarbons, preferably dichloromethane, trichloromethane, tetrachloromethane, trichloroethane and tetrachloroethane, in particular dichloromethane.

The reaction is preferably carried out by contacting a solution of the compound of formula (II) in a chlorinated solvent, with "stabilised" nitric acid dispersed in the same solvent. The solution of the compound of formula (II) should be as homogeneous as possible. For example, a homogeneous solution of 1,4-butanediol in dichloromethane can be obtained at concentrations ranging from about 60 to about 75%, preferably from about 65 to about 70% w/w.

The weight ratio of "stabilised" nitric acid to compounds of formula (II) ranges from about 10 : 1 to about 15 : 1, depending on the characteristics of the compound of formula (II).

For example, if the compound of formula (II) is 1,4-butanediol, the weight ratio of "stabilised" nitric acid to 1,4-butanediol preferably ranges from about 11 : 1 to about 14.5 : 1. Thus, the concentration of 1,4-butanediol

in the nitration mixture ranges from about 2.1 to about 2.8% w/w, preferably from about 2.3 to about 2.6% w/w.

The mononitration of compounds of formula (II) is carried out at a reaction temperature lower than room temperature, preferably equal to or lower than 0°C, more preferably at 0°C, for reaction times ranging from about 10 to about 30 minutes, preferably from about 15 to about 20 minutes. The progress of the reaction can be monitored with conventional analytical methods and the optimal reaction time can thus be determined. The reaction is typically quenched with cold water (precooled to a temperature lower than about 6°C, preferably lower than about 3°C).

The process of the invention allows to obtain the compounds of formula (I) in advantageous yields from the industrial standpoint and is definitely less hazardous than nitration with concentrated nitric acid, and optionally sulfuric acid, in the presence of urea.

As far as yields are concerned, the mononitration of 1,4-butanediol according to the invention affords 1,4-butanediol mononitrate with molar yield ranging from about 30% to about 40%, with a selectivity, expressed as percentage ratio 1,4-butanediol mononitrate/(1,4-butanediol mononitrate + 1,4-butanediol dinitrate) equal to about 70-75%.

The nitration mixture, which is a further object of the invention, is a crude mixture dispersed in the organic chlorinated solvent comprising the compound of formula (I), in amount of about 2% w/w, the corresponding dinitrate by-product, the unreacted compound of formula (II) and unreacted nitric acid, and other by-products deriving from dehydration and/or oxidation. The nitration mixture is first partially neutralized with a concentrated sodium hydroxide solution. Most of the unreacted compound of formula (II) is extracted in the resulting aqueous sodium nitrate solution. The organic phase is then concentrated by evaporation of the solvent and neutralized with a

diluted sodium carbonate or sodium hydroxide solution. The mixture in said chlorinated organic solvent generally contains a compound of formula (I) in amount approx. ranging from 11% to 15% w/w, the corresponding dinitrated by-product in amount approx. ranging from 3 to 4, 5% w/w; and the corresponding diol of formula (II) in amount approx. ranging from 0.2 to 0.8% w/w [compared with the compound of formula (I)]. Mixtures with total nitroesters concentrations higher than 15% w/w have explosive characteristics and are potentially hazardous to operators. The compounds of formula (I) can be separated from the corresponding dinitrated by-products and from traces of the corresponding diols of formula (II) according to known methods.

The following examples further illustrate the invention.

EXAMPLES

Example 1 - preparation of "stabilised" nitric acid

A stainless steel reactor, equipped with condenser and stirrer, is loaded with 90 kg of diluted nitric acid (84.7%) having a nitrous acid content of about 0,09% and 675 g of urea beads is added under stirring. After about 90 minutes, removal of nitrous acid and nitrogen oxides is controlled by colour inspection and permanganate titration. If necessary, further urea is added in small portions.

Example 2 - 1,4-Butanediol-mononitrate

A stainless steel reactor is loaded in succession with 931 g of dichloromethane and 385 g of "stabilised" nitric acid. The dispersion is cooled to about 0°C under stirring, thereafter 50 g of 1,4-butanediol in dichloromethane (70/30) is added in a single portion. The reaction mixture is kept under stirring at a temperature ranging from about -2°C to 2°C. The kinetics of the nitration is monitored by taking samples of the mixture during the reaction. After 20 minutes the reaction is quickly quenched by pouring it

into a water/ice mixture (385 g), then neutralized with 433 g of 40% NaOH, keeping the temperature below 15°C. The organic phase (980 g), which contains 1,4-butanediol-mononitrate (19.2 g), 1,4-butanediol-dinitrate (6.4 g) and 1,4-butanediol (0.1 g), is separated and concentrated to 15%. The molar yield of 1,4-butanediol-mononitrate in the crude solution amounts to 36.6%. The solution is then subjected to the subsequent purification.

The following compounds can also be obtained through the same procedure: ethanediol-mononitrate, 1,3-propanediol-mononitrate, 1,5-pentanediol-mononitrate and 1,6-hexanediol-mononitrate.

Example 3 - 1,4-Butanediol-mononitrate

Following the procedure described in example 2, the reactor is loaded with 21.78 kg of dichloromethane and 9 kg of "stabilised" nitric acid. The mixture is cooled to -5-0°C, then 1.26 kg of a mixture of 1,4-butanediol (0.819 kg) in dichloromethane (0.441 kg) is added under vigorous stirring in a single portion. After 18 minutes, the reaction is rapidly quenched in water (9 kg), precooled at 2°C. The solution is then neutralized with 40% sodium hydroxide (9 kg). The organic phase (23 kg), which contains 1,4-butanediol-mononitrate (0.46 kg), 1,4-butanediol-dinitrate (0.155 kg) and 1,4-butanediol (0.002 kg), is separated and concentrated to 15%. The molar yield of 1,4-butanediol-mononitrate in the crude solution amounts to 37.4%. The solution is then subjected to the subsequent purification.

CLAIMS

1. A process for the preparation of a compound of formula



wherein A is a C₂-C₆ alkylene chain,

comprising the nitration of a compound of formula



wherein A is as defined above,

with "stabilised" nitric acid.

2. A process as claimed in claim 1, wherein the compound of formula (I) is ethanediol-mononitrate; 1,3-propanediol-mononitrate; 1,4-butanediol-mononitrate; 1,5-pentanediol-mononitrate or 1,6-hexanediol-mononitrate.

3. A process according to any one of claims 1 or 2, wherein the "stabilised" nitric acid has a concentration ranging from 83 to 85% and is substantially free from nitrous acid and nitrogen oxides.

4. A process according to any one of claims 1-3, wherein the reaction is carried out in a water-immiscible chlorinated organic solvent.

5. A process as claimed in claim 4, wherein the chlorinated organic solvent is a mono-, di-, tri- or tetra-chloro C₁-C₄-alkyl hydrocarbon.

6. A process according to any one of claims 1-5, wherein the weight ratio of "stabilised" nitric acid to the compound of formula (II) ranges from 10 : 1 to 15 : 1.

7. A process according to any one of claims 1-5, wherein the nitration is carried out for a time ranging from 10 to 30 minutes.

8. A process according to any one of claims 1-7, wherein the compound of formula (II) is 1,4-butanediol and the weight ratio of "stabilised" nitric acid to butanediol ranges from 11: 1 to 14.5: 1.

9. Nitration mixture in a water-immiscible organic chlorinated solvent

comprising a compound of formula (I), as obtainable by the process of claim 1.

10. "Stabilised" nitric acid characterized in that it has a concentration ranging from 83 to 85% and is substantially free from nitrous acid and nitrogen oxides.

11. Process for the preparation of "stabilised" nitric acid comprising the dilution of fuming nitric acid with water to a concentration of about 83 - 85% and treatment with urea or sulfamic acid, in amount ranging from 0.3 to 1% w/w, for a time ranging from 80 to 130 minutes.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 03/12376

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07C201/02 C07C203/04 C01B21/46

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07C C01B

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, BEILSTEIN Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	GB 1 040 139 A (NITROCHEMIE GMBH) 24 August 1966 (1966-08-24) the whole document	1-11
Y	DE 20 39 609 A (COUNCIL SCIENT IND RES) 17 February 1972 (1972-02-17) page 2, paragraph 4 -page 3, paragraphs 1,2	1-11
Y	US 5 162 568 A (DOUTY CHARLES F) 10 November 1992 (1992-11-10) abstract	1-11
Y	EP 1 038 862 A (DINAMITE DIPHARMA S P A) 27 September 2000 (2000-09-27) example 2	1-11



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

29 January 2004

Date of mailing of the international search report

12/02/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Bonnevalle, E

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 03/12376

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
GB 1040139	A	24-08-1966	DE 1160427 B BE 632627 A CH 420093 A FR 1357403 A	02-01-1964 15-09-1966 03-04-1964
DE 2039609	A	17-02-1972	DE 2039609 A1 CH 540878 A	17-02-1972 15-10-1973
US 5162568	A	10-11-1992	NONE	
EP 1038862	A	27-09-2000	IT MI990627 A1 EP 1038862 A2	26-09-2000 27-09-2000